

CRONO[®] Motion & Energy



Data Sheet

Highly Bioavailable Natural Omega-3 Supplement Rich in DHA/EPA and Vitamins A and D₃, Formulated to Optimize the Musculoskeletal System, Cardiovascular Function and Athletic Performance

Active ingredients (per ml)*:

Omega-3	283,2 mg
DHA (21-23%)	190,8 mg
EPA (5-7%)	51,0 mg
Other Omega 3	41,4 mg
Vitamin A	1.737 UI
Vitamin D ₃	34 UI

*Variable quantity because of being a natural product

Composition: Cold extracted fish oil rich in Omega 3, vitamins A and D₃.

Analytical constituents: Crude oils and fats 99.54%.

Mechanism of action:

- Both fats as well as oils are made up of a mixture of fatty acids. Polyunsaturated fatty acids are also known as Essential Fatty Acids (EFA) because they cannot be synthesized by the body and must, therefore, be supplied through the diet.
 - There are two types of EFAs: Omega-3 (ALA, DHA, EPA...) and Omega-6 (LA, AA...). Cold-water fish are the richest natural source of Omega-3, particularly DHA and EPA. There are sources from plants such as linseed oil, but these are rich in ALA. Dietary ALA must be transformed by the body into biologically active Omega-3 DHA and EPA, however, this bioconversion is very inefficient: in people ALA conversion to DHA and EPA is below 0.10% and 10% respectively (Williams and Burdge, 2006). It has been shown that in horses supplemented with equivalent quantities of Omega-3 from linseed or fish oil, the Omega-3 plasma content was increased only in horses supplemented with fish oil (Siciliano et al., 2003). Other studies in horses have proven that fish oil significantly increased the concentration of DHA and EPA in plasma (Hall et al., 2004b; O'Connor et al., 2007; King et al., 2008).
 - Omega-6 has been called "bad Omega" because the eicosanoids (prostaglandins, thromboxanes and leukotrienes) derived therein show pro-inflammatory activity, while the activity of Omega-3 is anti-inflammatory. This does not imply that Omega-6 should be eliminated from the diet, but there should be an appropriate ratio between the two. In modern human and animal diets this proportion is very biased towards Omega-6 and this imbalance may favour certain pathologies such as osteoarthritis, atherosclerosis, asthma, cancer, neurological and behavioural disorders. On increasing Omega-3 intake, EPA competes as a substrate for enzymes in the arachidonic acid cascade (AA, an Omega-6 precursor of the destructive eicosanoids) and this abundance of Omega-3 can replace the destructive subtype AA products in favour of the protective subtypes (Robinson and Stone, 2006).
- **DHA/EPA in the Musculoskeletal System:**
Omega-3 reduces inflammation of the joints and helps heal and restore connective tissues damaged by stress from exercise.
In vitro studies have proven that DHA and EPA increase collagen synthesis and decrease PGE2 inflammation mediators.
It has been shown that supplementation with DHA/EPA in horses diagnosed with arthritis increases stride length, indicating a possible reduction in pain (Woodward et al., 2005). DHA and EPA have been tested with considerable success (especially in the early stages of illness) in treating the symptoms of rheumatoid arthritis in people, and although they probably do not slow the progression of the disease itself, they very satisfactorily moderate the symptoms. A recent study in people has shown that



Features

Reduces pain and inflammation in joints, tendons and muscles.

Improves cardiovascular function - Optimises athletic performance.

Fast and safe energy source - Improves response to prolonged exertion and delays the onset of fatigue.

Very safe - Ideal for extended treatment.

Does not contain doping substances.

Patented Omega-3 cold extraction method - Maximum bioactivity.

Environmentally friendly.



VetNova

CRONO[®] Motion & Energy



Highly Bioavailable Natural Omega-3 Supplement Rich in DHA/EPA and Vitamins A and D₃, Formulated to Optimize the Musculoskeletal System, Cardiovascular Function and Athletic Performance

the combined use of DHA/EPA and glucosamine greatly improves the comfort of patients with osteoarthritis of the knee and hip.

• DHA/EPA and Athletic Performance:

Different trials show that Omega-3 reduces blood viscosity. This decreases the heart rate and increases oxygen flow to the muscles, thus improving performance and response to stress. This feature of Omega-3 is particularly valuable for endurance horses. In a study in horses, heart rate after a stress test was lower in the group supplemented with DHA/EPA than in the control group while the lactate values were similar in the two groups, suggesting that horses supplemented DHA/EPA will take longer to become fatigued. This cardioprotective effect has been shown extensively in humans.

Studies in rodents suggest that EPA prevents muscular damage by inhibiting the cyclooxygenase pathway (Jackson et al., 1988; Tisdale, 1996).

Competition horses largely depend on anaerobic glycolysis for energy, while fatigue is more likely to be associated with the accumulation of lactic acid. DHA/EPA reduce this accumulation and at the same time balance the increase in production of enzymes and hormones related to the competition, thus increasing speed and endurance. Omega-3 increases tissue elasticity and may help reduce the incidence of EIPH (Exercise Induced Pulmonary Haemorrhage).

• DHA/EPA as a Source of Energy:

Carbohydrate reserves (as glycogen in the liver and muscles) are used only to maintain short-term exertion (e.g. sprinting), but they do not supply enough energy alone for long-term exertion (e.g. endurance); when they are depleted premature fatigue arises. Additionally, horses requiring a high-energy diet (sport horses or those with poor body condition) are at high risk of colic and/or laminitis associated with excessive fermentation of a high carbohydrate diet (e.g. grain).

Omega-3 is a healthy source of highly concentrated (providing 2.5 times more energy than an equivalent quantity of grain or feed), digestible and easy to metabolise fat energy, and therefore, it is a very safe alternative to carbohydrates. In the body, dietary fat is directly used as energy or stored in adipose tissue, under the skin or around the organs, improving body condition. A smaller quantity of fat is stored in the muscle to be used as energy for muscular work, improving athletic performance, although an adaptation period of around three weeks is needed for the horse's metabolism to achieve an increase in the use of fat as fuel for exercise and to reduce the expense of liver and muscular glycogen.

• DHA/EPA in Reproduction and Breeding:


In stallions with low fertility, DHA improves the characteristics of the spermatozoa, the concentration in the seminal fluid and the daily spermatozoa production. Other studies in horses have shown an improvement in the viability of frozen and refrigerated seminal fluid in stallions supplemented with DHA/EPA.

DHA is vital for the cerebral, cognitive and visual development of the foetus and the foal. Administration of Omega-3 during pregnancy contributes to optimal development of the foetus, preventing defects. Additionally, mares incorporate Omega-3 into the colostrum and milk, and some studies suggest that the foals of mares supplemented with Omega-3 appear to have a stronger immune system. It has been proven in dogs that the offspring of mothers with high levels of DHA are more intelligent and easier to train (Heinemann and Bauer, 2006; Zicker et al., K.M., 2012).

There is testimonial evidence which suggests that Omega-3 improves the regeneration of reproductive organs in mares after birth, thus facilitating future conception. It has also been shown that DHA/EPA give rise to faster and more consistent cycles. Other findings suggest that mares with a history of abortion may benefit from nutritional support with Omega-3.

• Other Benefits of DHA/EPA:

- Skin/Coat: EFAs are useful for managing allergic dermatological conditions such as summer dermatitis and they improve the external appearance of the horse (skin and coat).



CRONO[®] Motion & Energy



Highly Bioavailable Natural Omega-3 Supplement Rich in DHA/EPA and Vitamins A and D₃, Formulated to Optimize the Musculoskeletal System, Cardiovascular Function and Athletic Performance

Data Sheet



- Due to their anti-inflammatory and immunopotentiating action, DHA and EPA may help in respiratory conditions such as RAO (Recurrent Airway Obstruction), IAD (Airway inflammation), laminitis, etc.
- Renal: Various studies have proven that EFAs reduce the progression of chronic renal disease, particularly at the glomerular level, reducing the proteinuria.
- In other animal species, Omega-3 has beneficial effects such as reducing blood triglycerides and increasing membrane fluidity and insulin sensitivity (Mueller and Talbert, 1988; Simopoulos, 1991).
- Nervous System: Research trials for Alzheimer's disease in rodents have proven that DHA has anti-inflammatory properties and reduces amyloid levels and plaque formation (Horrocks and Yeo, 1999; Lim et al., 2003) suggesting that Omega-3 can contribute to preserving the physical, mental and sensory activity of older horses.

• Benefits of Vitamins A and D₃:

Vitamin A contributes to normal iron metabolism, maintaining natural protective barriers such as the skin and mucous membranes, the normal functioning of the immune system and, together with DHA, proper maintenance of vision.

Vitamin D₃ is essential for the normal use and absorption of calcium and phosphorus, it contributes to bone, dental and muscular health and to the optimal functioning of the immune system.

Indications: Recommended by veterinarians to optimise the functionality of the musculoskeletal system (joints, tendons and muscles), and to improve the cardiovascular function and athletic performance of horses under joint/muscular stress due to intense exercise or competition. In older horses with age-related joint degeneration, CRONO[®] Motion & Energy helps restore mobility and quality of life for the animal.

Also recommended for horses that require a high-energy diet (sport horses or those with poor body condition) as a source of digestible, easy to metabolise and very safe concentrated energy, in order to avoid metabolic disorders associated with diets rich in starch.

Improves the quantity and quality of spermatozoa in stallions, supports pregnancy and lactation in mares and facilitates the cerebral and visual development of the foetus and nursing foal.

Target species: Equidae.


Dosage and administration: Shake before use. Administer mixed with feed or orally through a syringe. Daily dose: Initial (1 month), 30 ml; Maintenance (from the 2nd month), 15 ml. Dose for an average size adult horse (500 kg), adjust the dose based on weight. To improve initial acceptance we recommend dividing the dose into 2 separate doses: morning and evening. The effects of CRONO[®] Motion & Energy start to be seen after 3-4 weeks of application.

Safety: CRONO[®] Motion & Energy has no contraindications and can be taken long-term or for life.

Manufacture and the Environment:

- The Omega-3 in the CRONO[®] Motion & Energy formula is obtained through an exclusive patented method; 100% natural cold extraction which keeps the DHA and EPA molecular structures intact, thus ensuring their maximum bioavailability and bioactivity. Other products use boiling processes at high temperatures or chemical solvents for extracting fish oil.
- CRONO[®] Motion & Energy is 100% environmentally friendly as it does not generate waste, making good use of all the raw material used.
- CRONO[®] Motion & Energy is manufactured in a pharmaceutical grade plant in Galicia, Spain.

VetNova



CRONO[®] Motion & Energy



Data Sheet


Highly Bioavailable Natural Omega-3 Supplement Rich in DHA/EPA and Vitamins A and D₃, Formulated to Optimize the Musculoskeletal System, Cardiovascular Function and Athletic Performance

Warnings: This product does not contain any substances which are banned in competition. Keep the container tightly closed in a cool, dry place, away from direct sunlight and out of reach of children and animals. Liquid at room temperature, it may form precipitates if refrigerated. Due to being an oily product it may experience minor expansions and/or contractions during transport, causing minor losses or deformations of the container. Raw material for feed.

Presentation: 930 ml (31-62 doses for 1-2 months' supply).

Bibliography:

- Agostoni, C., Trojan, S., Bellu, R., Riva, E., Bruzzese, M.G., Giovannini, M. Developmental quotient at 24 months and fatty acid composition of diet in early infancy: follow up study. Arch Dis Chil 1997; 76:421-424.
- Aguado Acín, M.P. Vitamina D y osteoporosis. ¿Es suficiente tomar el sol? Rev Esp Reumatol 2001; 28:101-105.
- Angerer, P., Von Schacky, C. Ω -3 polyunsaturated fatty acids and the cardiovascular system. Curr Opin Lipidol 2000; 11:57-63.
- Bahadori, B., Uitz, E., Thonhofer, R., Trummer, M., Pestemer-Lanch, I., McCarty, M., Krejs, G.J. Omega-3 fatty acids infusions as adjuvant therapy in rheumatoid arthritis. JPEN J Parenter Enteral Nutr 2010; 34:151-155.
- Bazan, N.G. Cell survival matters: docosahexaenoic acid signaling, neuroprotection and photoreceptors. Trends Neurosci 2006; 29:263-271.
- Burr, G.O., Burr, M.M. On the nature and role of fatty acids essential in nutrition. J Biol Chem 1930; 86: 587-621.
- Calder, P.C. Polyunsaturated fatty acids, inflammatory processes and autoimmune diseases. Mol Nutr Food Res 2008; 52(8):885-897.
- Calon, F., Cole, G. Neuroprotective action of omega-3 polyunsaturated fatty acids against neurodegenerative diseases: evidence from animal studies. Prostaglandins Leukot Essent Fatty Acids 2007; 77: 287-293.
- Cleland, L.G., French, J., Betts, H.W., Murphy, G.A., Elliott, M.J. Clinical and biochemical effects of dietary fish oil supplements in rheumatoid arthritis. J Rheumatol 1988;15:1471-1475.
- Davison, K.E. Should Horses Eat Fish? - Current Information on Omega-3 Supplements for Horses. Purina Mills, LLC
- Fritsch, D.A., T.A. Allen, C.E. Dodd, D.E. Jewell, K.A., Sixby, P.S. Leventhal, J. Brejda, Hahn, K.A. A multicenter study of the effect of dietary supplementation with fish oil omega-3 fatty acids on carprofen dosage in dogs with osteoarthritis. JAVMA 2010; 236:535-539.
- Gil-Campos, M., Dalmau Serra, J., Comité de Nutrición de la Asociación Española de Pediatría. Importancia del ácido docosahexaenoico (DHA): funciones y recomendaciones para su ingesta en la infancia. An Pediatr (Barc) 2010; 73(3):142.e1-142.e8.
- Guadalix, S., Jódar, E. Vitamina D y función muscular. Rev Esp Enferm Metab Oseas. 2007; 16(2):41-44.
- Hawkins Carranza, F. La vitamina D3 y el hueso. Rev Esp Enferm Metab Oseas. 2007; 16(3):45-47.
- Heinemann, K.M., Bauer, J.E. Docosahexaenoic acid and neurologic development in animals. J Am Vet Med Assoc. 2006 Mar 1; 228(5):700-5, 655.
- Horrocks, L.A., Farooqui, A.A. Docosahexaenoic acid in the diet: its importance in maintenance and restoration of neural membrane function. Prostaglandins Leukot Essent Fatty Acids 2004; 70: 361-372.
- Horrocks, L.A., Yeo, Y.K. Health benefits of docosahexaenoic acid (DHA). Pharmacol Res. 1999 Sep; 40(3):211-25.
- Hurst, S., Zainal, Z., Caterson, B., Hughes, C.E., Harwood, J.L. Dietary fatty acids and arthritis. Prostaglandins Leukot Essent Fatty Acids 2010; 82:315-318.
- Innis, S. Dietary (n-3) fatty acids and brain development. J Nutr 2007; 137:855-859.
- Jackson, M.J., Roberts, J., Edwards, R.H. Effects of dietary-fish-oil feeding on muscle growth and damage in the rat. Br J Nutr. 1988 Sep; 60(2):217-24.
- Jerosch, J. Effects of Glucosamine and Chondroitin Sulfate on Cartilage Metabolism in OA: Outlook on Other Nutrient Partners Especially Omega-3 Fatty Acids. Hindawi Publishing Corporation. Int J Rheumatol. 2011;
- Kalminj, S., Van Boxtel, M.P.J., Ocke, M., Verschuren, W.M.M., Kromhout, D., Launer, L.J. Dietary intake of fatty acids and fish in relation to cognitive performance at middle age. Neurology 2004 Jan 27; 62(2):275-80.
- King, S.S., AbuGhazaleh, A.A., Webel, S.K., Jones, K.L. Circulating fatty acid profiles in response to three levels of dietary omega-3 fatty acid supplementation in horses. J Anim Sci. 2008; 86:1114-1123.
- Lim, S.Y., Hoshiba, J., Moriguchi, T., Salem, N. Jr. N-3 fatty acid deficiency induced by a modified artificial rearing method leads to poorer performance in spatial learning tasks. Pediatr Res. 2005 Oct; 58(4):741-8.






Highly Bioavailable Natural Omega-3 Supplement Rich in DHA/EPA and Vitamins A and D₃, Formulated to Optimize the Musculoskeletal System, Cardiovascular Function and Athletic Performance

Data Sheet

- Lim, S.Y., Hoshiba, J., Salem, N. Jr. An extraordinary degree of structural specificity is required in neural phospholipids for optimal brain function: n-6 docosapentaenoic acid substitution for docosahexaenoic acid leads to a loss in spatial task performance. *J Neurochem.* 2005 Nov; 95(3):848-57.
- Lim, S.Y., Suzuki, H. Dose-response effect of egg-phosphatidylcholine on maze-learning ability and fatty acid composition of plasma and brain in aged mice fed an n-3 fatty acid-deficient diet. *Ann Nutr Metab.* 2002; 46(5):215-21.
- Lim, S.Y., Suzuki, H. Dose-response effect of docosahexaenoic acid ethyl ester on maze behavior and brain fatty acid composition in adult mice. *Int J Vitam Nutr Res.* 2002 Mar; 72(2):77-84.
- Lim, S.Y., Suzuki, H. Changes in maze behavior of mice occur after sufficient accumulation of docosahexaenoic acid in brain. *J Nutr.* 2001 Feb; 131(2):319-24.
- Lim, S.Y., Suzuki, H. Effect of dietary docosahexaenoic acid and phosphatidylcholine on maze behavior and fatty acid composition of plasma and brain lipids in mice. *Int J Vitam Nutr Res.* 2000 Sep;70(5):251-9
- Lim, S.Y., Suzuki, H. Intakes of dietary docosahexaenoic acid ethyl ester and egg phosphatidylcholine improve maze-learning ability in young and old mice. *J Nutr.* 2000 Jun; 130(6):1629-32.
- Manhart, D.R., Scott, B.D., Gibbs, P.G., Coverdale, J.A., Miller, E.M., Honnas, C.M., Hood, D.M. Markers of inflammation in arthritic horses fed omega-3 fatty acids. *Professional Animal Scientist* 2009; 25:155-160.
- Mayurasakom, K., Williams, J.J., Ten, V.S., Deckelbaum, R.J. Docosahexaenoic acid: brain accretion and roles in neuroprotection after brain hypoxia and ischemia. *Curr Opin Clin Nutr Metab Care* 2011; 14: 158-167.
- McCann, J.C., Ames, B.N. Is Docosahexaenoic acid, an n-3 long-chain polyunsaturated fatty acid, required for development of normal brain function? An overview of evidence from cognitive and behavioral tests in humans and animals. *Am J Clin Nutr* 2005; 82:281-295.
- Mesa García, M.D., Aguilera García, C.M., Gil Hernández, A. Importancia de los lípidos en el tratamiento nutricional de las patologías de base inflamatoria. *Nutr Hosp.* 2006; 21(2): 30-43.
- Morris, M.C., Evans, D.A., Bienias, J.L. et al. Consumption of fish and n-3 fatty acids and risk of incident Alzheimer disease. *Arch Neurol* 2003; 60: 940-946.
- O'Connor, C.I., Lawrence, L.M., St. Lawrence, A.C., Janicki, K.M., Warren, L.K., Hayes, S. The effect of dietary fish oil supplementation on exercising horses. *J Anim Sci.* 2004; 82:2978-2984.
- Lama, R.A., Moráis, A. Las grasas en la alimentación infantil. Importancia de los ácidos grasos poliinsaturados. *An Pediatr, Monogr* 2005; 3(1):16-23.
- Rueda, F., Domingo, J.C., Mach, N. Efectos de los ácidos grasos omega 3 y otros suplementos alimenticios en procesos patológicos relacionados con la tercera edad. *Rev Esp Nutr Hum Diet.* 2011; 15(1):20-29.
- Sánchez, A., Puche, R., Zeni, S., Oliveri, B., Galich, A.M., Meffi, L., Plantalech, L., Poudes, G., Bregni, C. Papel del calcio y de la vitamina D en la salud ósea (Parte I). *Rev Esp Enferm Metab Oseas.* 2002; 11(6):201-217.
- Sanhueza, J., Nieto, S., Valenzuela, A. Ácido docosahexaenoico (DHA), desarrollo cerebral, memoria y aprendizaje: la importancia de la suplementación perinatal. *Rev Chil Nutr* 2004; 31(2):84-92.
- Simopoulos, A.P. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed Pharmacother.* 2002; 56: 365-379.
- Sinn, N., Bryan, D. Effect of Supplementation with polyunsaturated fatty acids and micronutrients on learning and behavior problems associated with child ADHD. *J Dev Behav Pediatr* 2007; 28:82-91.
- Soderberg, M., Edlund, C., Kristensson, K., Dallner, G. Fatty acid composition of brain phospholipids in aging and in Alzheimer's disease. *Lipids* 1991; 26: 421-425.
- Studzinski, C.M., Araujo, J.A., Milgram, N.W. The canine model of human cognitive aging and dementia: pharmacological validity of the model for assessment of human cognitive-enhancing drugs. *Prog Neuropsychopharmacol Biol Psychiatry.* 2005 Mar; 29(3):489-98.
- Sontrop, L., Campbell, M.K. Omega-3 polyunsaturated fatty acids and depression: a review of the evidence and a methodological critique. *Prev Med* 2006; 42(1): 4-13.
- Stevens, L., Zentall, S., Deck, J., Abate, M., Watkins, B., Lipp, S. Essential fatty acid metabolism in boys with attention-deficit hyperactivity disorder. *Am J Clin Nutr* 1995; 62:761-768.
- Tanskanen, A., Hibbeln, J.R., Tuomilehto, J., Uutela, A., Haukkala, A., Viinamäki, H., Lehtonen, J., Vartiainen, E. Fish consumption and depressive symptoms in the general population in Finland. *Psychiatr Serv* 2001; 52(4): 529-531.
- Tisdale, M.J. Inhibition of lipolysis and muscle protein degradation by EPA in cancer cachexia. *Nutrition.* 1996 Jan;12(1 Suppl):S31-3
- Trebble, T.M., Wooten, S.A., Miles, E.A., Mullee, M., Arden, N.K., Ballinger, A.B., et al. Prostaglandin E2 production and T cell function after fish-oil supplementation: response to antioxidant co-supplementation. *Am J Clin Nutr* 2003; 78:376-382.



CRONO[®] Motion & Energy

Highly Bioavailable Natural Omega-3 Supplement Rich in DHA/EPA and Vitamins A and D₃, Formulated to Optimize the Musculoskeletal System, Cardiovascular Function and Athletic Performance

- Tully, A.M., Roche, H.M., Doyle, R., Fallon, C., Bruce, I., Lawlor, B., et al. Low serum cholesteryl esterdocosahexaenoic acid levels in Alzheimer's disease: a case-control study. *Br J Nutr* 2003; 89: 483-489.
- Valenzuela, A., Nieto, S. Ácido docosahexaenoico (DHA) en el desarrollo fetal y en la nutrición materno-infantil. *Rev Med Chil* 2001; 129:1203-1211.
- Valenzuela, A., Nieto, S. Ácidos grasos omega-6 y omega-3 en la nutrición perinatal: su importancia en el desarrollo del sistema nervioso y visual. *Rev Chil Pediatr* 2003; 74(2):149-157.
- Valenzuela, A., Sanhueza, J., Nieto, S. Docosahexaenoic acid (DHA), essentiality and requirements: why and how to provide supplementation. *Grasas & Aceites* 2006; 57:229-237.
- Valenzuela, A., Sanhueza, J. Aceites de origen marino; su importancia en la nutrición y en la ciencia de alimentos. *Rev Chil Nutr* 2009; 36:246-257.
- Valenzuela, R., Tapia, G., González, M., Valenzuela, A. Ácidos grasos omega-3 (EPA y DHA) y su aplicación en diversas situaciones clínicas. *Rev Chil Nutr* 2011; 3:356-367.
- Vineyard, K.R., Warren, L.K., Kivipelto, J. Effect of dietary omega-3 fatty acid source on plasma and red blood cell membrane composition and immune function in yearling horses. *J Anim Sci.* 2010; 88:248-257.
- Zicker, S.C., Jewell, D.E., Yamka, R.M., Milgram, N.W. Evaluation of cognitive learning, memory, psychomotor, immunologic, and retinal functions in healthy puppies fed foods fortified with docosahexaenoic acid-rich fish oil from 8 to 52 weeks of age. *J Am Vet Med Assoc.* 2012 Sep 1; 241(5):583-94. doi: 10.2460/javma.241.5.583.

If you are interested in any of the articles listed, please do not hesitate to request them through the following contacts: vetnova@vetnova.net, +34 918 440 273, or your VetNova or Distributor Sales Representative.



Data Sheet



VetNova

Teléf.: +34 918 440 273 · vetnova@vetnova.net · www.vetnova.net

VN-PUB-0079EN

