

CUTANIA® Chews

CUTANIA® MINI Chews



Next Generation Dermatological Supplement for Long Term Management of Hair and Skin Conditions in Dogs and Cats.

Data Sheet

Active Ingredients (per chew):

	CUTANIA®	CUTANIA® MINI	u.m.
<i>Perilla frutescens</i>	20	10	mg
<i>Vitis vinifera</i>	30	15	mg
MSM	75	37.5	mg
Zinc	5	2.5	mg
DHA (Omega-3)	7.5	3.75	mg
EPA (Omega-3)	11	5.5	mg
GLA (Omega-6)	15	7.5	mg
Hyaluronic Acid	5	2.5	mg
L-Cysteine	50	25	mg
DMG	50	25	mg
DL-Methionine	50	25	mg
L-Proline	25	12.5	mg
Vitamin A	37.5	18.75	IU
Vitamin B ₃	4	2	mg
Vitamin C	25	12.5	mg
Vitamin E	10	5	IU



Management of atopy and other chronic skin disorders.



Features

16 active ingredients having synergistic effect:

- *Perilla frutescens* (rich in polyphenols with antihistamine action)
- Omega-3 y Omega-6
- MSM
- Zinc
- Hyaluronic Acid
- Vitamins A, B₃, C and E
- Cysteine, Methionine and Proline

Composition (in descending order): Corn starch, brewer's yeast, imitation chicken vegetable flavouring, water, glycerine, soya lecithin, black currant oil, oatmeal, barley, rice and rye, safflower oil, oat bran, methylsulphonylmethane (MSM), vitamin C, L-cysteine, N, N-Dimethylglycine HCl (DMG), DL-Methionine, maltodextrin mixture, sodium alginate and calcium sulphate, fish oil, *Vitis vinifera* extract, L-Proline, *Perilla frutescens* seed extract, zinc, natural vitamin E, sorbic acid, low molecular weight hyaluronic acid (HyaMax®), niacinamide, propionic acid, mixture of natural tocopherols combined with small quantities of silicon dioxide, citric acid, vegetable oil and rosemary extract, vitamin A (retinyl acetate).

Analytical Constituents (CUTANIA®/CUTANIA® Mini): Crude protein 17.1/26.8%; crude oils and fats 13.6/10.33%; crude fibre 0/1.9%; crude ash 4.07/7.46%; moisture 14.3/6.28%; DHA 0.21/0.25%; EPA 0.31/0.36%; GLA 0.43/0.49%.

Essential Nutritional Characteristics: Contains essential fatty acids omega-3 and 6.

Properties:

CUTANIA® and CUTANIA® Mini contain a complete and unique combination of active ingredients that act synergistically to promote the improvement of skin and hair conditions and stimulate natural skin repair.

- ***Perilla frutescens*** is a very rich source of polyunsaturated fatty acids. Compared to other vegetable oils, it contains the highest proportion of Omega-3 (ALA) fatty acids, high percentages of Omega-6, such as Linoleic Acid, with excellent antihistamine and antioxidant properties and Omega-9 (Oleic Acid). Furthermore, it is a rich source of phenolic compounds, flavonoids and anthocyanins such as rosmarinic acid and luteolin. These compounds are known for their antioxidant properties, their anti-microbial, anti-inflammatory and especially anti-histaminic action.
- **Omega-3 and 6** fatty acids are associated with benefits in a wide range of inflammatory processes, asthma, allergies, anti-tumour, etc. Supplementation with polyunsaturated fatty acids improves, and sometimes eliminates, the itching in allergic dermatopathies, such as atopy and allergies to fleabites in the case of dogs, and allergic and idiopathic itching in cats. Numerous studies have shown that they suppress a wide range of allergic mediators in allergic and inflammatory responses, due to the effects of Alpha-Linolenic Acid (ALA) on leukotrienes. Moreover, they improve skin elasticity.
- **MSM** (Methylsulphonylmethane) has analgesic and anti-inflammatory properties. MSM reduces scar tissue, speeding up wound healing and it promotes collagen production. In healthy dogs MSM improves overall health and promotes growth and luster of the coat.

Dual Action:

- Breaks the vicious cycle
Itching-Inflammation-Discomfort»»
Scratching-Licking-Self-Harm....
- Stimulates keratin and collagen production, activating the natural repair mechanisms in the skin.

Promotes fast and healthy healing.

Alternative to corticosteroids for controlling clinical symptoms after overcoming the acute phase.

Supportive nutraceutical for follicular dysplasias.

Highly palatable chews.

CUTANIA® Chews

CUTANIA® MINI Chews

Next Generation Dermatological Supplement for Long Term Management of Hair and Skin Conditions in Dogs and Cats.



Data Sheet

- **Hyaluronic Acid** is responsible for initiating the recovery and stabilisation process when the skin suffers injury by inducing the activation of receptors and immune system cells. In addition to its healing properties, it also acts as an antimicrobial barrier. It has the ability to retain large quantities of water, useful for lubricating and stimulating collagen production.
- **DMG** (DiMethylGlycine) improves the immune system and also has anti-inflammatory and antioxidant properties.
- **Vitamins A** (Retinol), **B₃** (Niacin/Niacinamide), **C** and **E** (Alpha-Tocopherol) are necessary for the skin's proper healing process, being important for their role in the formation and maintenance of collagen, as well as for their antioxidant effect that helps control the action of free radicals, essential for the prevention of cell ageing.
Moreover, Vitamins B₃ and C reduce skin hyperpigmentation.
- **Zinc** is a mineral which is involved in the immune response, necessary for the formation of proteins and collagen. It works synergistically with Vitamin E. At the cutaneous level it actively takes part in the regeneration of the extra-cellular matrix, healing processes, connective tissue repair, inflammation, and cell growth.
- **L-Cysteine** is a constituent of alpha-keratin, the most abundant protein in the skin, hair and nails. It takes part in the healing process.
- **DL-Methionine** and **L-Proline** are involved in collagen production and in wound repair. They have antioxidant properties and are also involved in immune responses. Methionine also reduces the release of histamine.
- **Vitis vinifera** (grape seed extract) contains proanthocyanidins, which improve the reconstruction of connective tissue, reduce the damage caused by prostaglandins, inhibit the release of histamine and protect collagen and elastin fibres through their antioxidant action.

Mechanism of Action:

- **Phase I:** *Perilla frutescens*, MSM, Omega-3, Omega-6, L-Methionine, *Vitis vinifera* and DMG reduce itching, discomfort and inflammation, minimising the response of the pet (scratching, licking, self-harm) which perpetuates the process.
- **Phase II:** Hyaluronic Acid, MSM, essential amino acids Cysteine, Methionine and Proline, Vitamins A, B₃, C and E, Zinc and *Vitis vinifera* stimulate collagen production and skin hydration, speeding up recovery and healing.

Indications: Indicated in hair and skin conditions to break the vicious cycle *Itching/Inflammation/Discomfort*»*Scratching/Licking/Self-harm*..., and to stimulate collagen and keratin production in order to activate the skin's natural repair mechanisms. Essential in long term treatments because it is able to maintain appropriate conditions for recovery of wounds and the proper functioning of epithelial tissue. CUTANIA® is a good alternative to corticosteroids to control the clinical conditions once the acute phase has passed.

- Atopic dermatitis: environmental, seasonal allergies...
- Food allergies.
- Skin traumas: post-surgical healing, burns, etc.
- Infectious (*Malassezia*, bacteria or fungi) and parasitic (scabies, fleas...) dermatitis.
- Ichthyosis, dry, moist seborrhoeic dermatitis...
- Localised dermatitis: moist eczema, pododermatitis, licked granuloma, etc.
- Follicular dysplasia...

Target Species: Dogs and cats.



VetNova

CUTANIA® Chews

CUTANIA® MINI Chews



Next Generation Dermatological Supplement for Long Term Management of Hair and Skin Conditions in Dogs and Cats.

Data Sheet

Directions of Use:

CUTANIA® Chews: 1 chew/7.5 kg.

CUTANIA® Mini Chews: 1 chew/3.75 kg.

Do not administer to animals with diarrhoea.

Warnings: VetNova is a pioneer in the development of Chews technology to facilitate administration of supplements to cats and dogs. Unlike tablets, capsules, etc. which are administered in a "forced" manner to the mouth to ensure the product is taken, Chews should be administered freely in the pet's bowl and the pet left to take them voluntarily. Some shy cats and dogs may need a longer time to fully accept them, but once they do the daily intake is easier and more satisfactory. The following strategies can be used during the first week to facilitate initial acceptance: 1) Reduce the dose and gradually increase it, 2) Divide the daily dose into two separate doses (morning and evening), 3) Crush the Chew and mix it with pate or any food which the pet enjoys, etc.

Presentation:

- CUTANIA® Chews: 60 Chews (1 month for a 15 kg dog).
- CUTANIA® Mini Chews: 60 Chews (2 months for a cat or miniature dog).

Bibliography:

- Asif M. Health effects of omega-3,6,9 fatty acids: Perilla frutescens is a good example of plant oils. *Orient Pharm Exp Me.* 2011 Mar; 11 (1): 51-59.
- Araujo ML, Burgos MG, Moura IS. Nutritional influences in psoriasis. *Universidade Federal de Pernambuco, Recife, PE, Brazil. An Bras Dermatol.* 2009 Jan-Feb;84(1):90-2.
- Aust MC, Fernandes D, Kolokythas P, Kaplan HM, Vogt PM. Percutaneous collagen induction therapy: an alternative treatment for scars, wrinkles, and skin laxity. *Klinik für Plastische, Hand- und Wiederherstellungschirurgie, Medizinische Hochschule Hannover, Hannover, Germany. Plast Reconstr Surg.* 2008 Apr;121(4):1421-9.
- Baumann LS. Less-known botanical cosmeceuticals. *Division of Cosmetic Dermatology, University of Miami School of Medicine, Miami, Florida 33125, USA. Dermatol Ther.* 2007 Sep-Oct;20(5):330-42.
- Berardesca E, Cameli N, Cavallotti C, Levy JL, Piérard GE, de Paoli Ambrosi G. Combined effects of silymarin and methylsulfonylmethane in the management of rosacea: clinical and instrumental evaluation. *San Gallicano Dermatological Institute, Rome, Italy. J Cosmet Dermatol.* 2008 Mar;7(1):8-14.
- Brien S, Prescott P, Bashir N, Lewith H, Lewith G. Systematic review of the nutritional supplements dimethyl sulfoxide . (DMSO) and methylsulfonylmethane (MSM) in the treatment of osteoarthritis. *Osteoarthritis Cartilage.* 2008 Nov;16(11):1277-88. doi: 10.1016/j.joca.2008.03.002.
- Calder PC, Albers R, Antoine JM, Blum S, Bourdet-Sicard R, Ferns GA, Folkerts G, Friedmann PS, Frost GS, Guarner F, Løvik M, Macfarlane S, Meyer PD, M'Rabet L, Serafini M, van Eden W, van Loo J, Vas Dias W, Vidry S, Winkhofer-Roob BM, Zhao J. Inflammatory disease processes and interactions with nutrition. *School of Medicine, University of Southampton, Southampton SO16 6YD, UK. Br J Nutr.* 2009 May;101 Suppl 1:S1-45.
- Colella G, Cannavale R, Vicidomini A, Rinaldi G, Compilato D, Campisi G. Efficacy of a spray compound containing a pool of collagen precursor synthetic aminoacids (l-proline, l-leucine, l-lysine and glycine) combined with sodium hyaluronate to manage chemo/radiotherapy-induced oral mucositis: preliminary data of an open trial. *Department of Head and Neck Surgery, Second University of Naples, Italy. Int J Immunopathol Pharmacol.* 2010 Jan-Mar;23(1):143-51.
- Damian DL. Photoprotective effects of nicotinamide. *Dermatology, Gloucester House Level 3, University of Sydney at Royal Prince Alfred Hospital Camperdown, NSW, 2050, Australia. J Drugs Dermatol.* 2010 Jun;9(6 Suppl):S72-81; quiz s82-3.
- Davis EC, Callender VD. Postinflammatory hyperpigmentation: a review of the epidemiology, clinical features, and treatment options in skin of color. *J Clin Aesthet Dermatol.* 2010 Jul;3(7):20-31.
- De Luis D, Aller R. Systematic review of nutritional support in pressure ulcer. *Sección de Endocrinología y Nutrición Clínica. Facultad de Medicina. Valladolid. An Med Interna.* 2007 Jul;24(7):342-5.
- Fleck CA. Managing ichthyosis: a case study. *Ostomy Wound Manage.* 2006 Apr;52(4):82-6, 88, 90.



VetNova

CUTANIA® Chews

CUTANIA® MINI Chews



Next Generation Dermatological Supplement for Long Term Management of Hair and Skin Conditions in Dogs and Cats.

Data Sheet

- Fowler JF Jr, Woolery-Lloyd H, Waldorf H, Saini R. Innovations in natural ingredients and their use in skin care. University of Louisville, Division of Dermatology, Louisville, KY 40202, USA. *Photochem Photobiol Sci.* 2010 Apr;9(4):578-85.
- Fu PP, Xia Q, Boudreau MD, Howard PC, Tolleson WH, Wamer WG. Physiological role of retinyl palmitate in the skin. National Center for Barrager E, Veltmann JR Jr, Schauss AG, Schiller. RNA multicentered, open-label trial on the safety and efficacy of methylsulfonylmethane in the treatment of seasonal allergic rhinitis. GENESIS Center for Integrative Medicine, Graham, WA, USA. *J Altern Complement Med.* 2002 Apr;8(2):167-73.
- Gannon J, Kendall R. A critical evaluation of N,N-dimethylglycine (DMG) and disopropylammonium dichloroacetate (DIPA) on the performance of racing greyhounds. In: Anonymous. 7 ed. 1982;9.
- Graber CD, Goust JM, Glassman AD, Kendall R, Loadholt CB. Immunomodulating properties of dimethylglycine in humans. *J Infect Dis.* 1981 Jan;143(1):101-5.
- Guo R, Pittler MH, Ernst E. Herbal medicines for the treatment of allergic rhinitis: a systematic review. Department of Complementary Medicine, Peninsula Medical School, University of Exeter, England. *An Allergy Asthma Immunol.* 2007 Dec;99(6):483-95.
- Jerajani HR, Mizoguchi H, Li J, Whittenbarger DJ, Marmor MJ. The effects of a daily facial lotion containing vitamins B3 and E and provitamin B5 on the facial skin of Indian women: a randomized, double-blind trial. Department of Dermatology, L.T.M. Medical College, L.T.M. General Hospital, Mumbai, India. *Indian J Dermatol Venereol Leprol.* 2010 Jan-Feb;76(1):20-6.
- Johansson S, Lönnqvist A, Ostman S, Sandberg AS, Wold AE. Long-chain polyunsaturated fatty acids are consumed during allergic inflammation and affect T helper type 1 (Th1)- and Th2-mediated hypersensitivity differently. Department of Chemical and Biological Engineering, Chalmers University of Technology, Gothenburg, Sweden. *Clin Exp Immunol.* 2010 Jun;160(3):411-9. Epub 2010 Feb 10.
- Jun JI, Lau LF. The matricellular protein CCN1 induces fibroblast senescence and restricts fibrosis in cutaneous wound healing. Department of Biochemistry and Molecular Genetics, University of Illinois at Chicago College of Medicine, Chicago, IL 60607-7170, USA. *Nat Cell Biol.* 2010 Jul;12(7):676-85. Epub 2010 Jun 6.
- Kim LS, Axelrod LJ, Howard P, Buratovich N, Waters RF. Efficacy of methylsulfonylmethane (MSM) in osteoarthritis pain of the knee: a pilot clinical trial. *Osteoarthritis Cartilage.* 2006 Mar;14(3):286-94. Epub 2005 Nov 23.
- Kim WS, Yang YJ, Min HG, Song MG, Lee JS, Park KY, Kim JJ, Sung JH, Choi JS, Cha HJ. Accelerated wound healing by S-methylmethionine sulfonium: evidence of dermal fibroblast activation via the ERK1/2 pathway. Department of Dermatology, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, Korea. *Pharmacology.* 2010;85(2):68-76. Epub 2010 Jan 21.
- Kim YH, Kim DH, Lim H, Baek DY, Shin HK, Kim JK. The anti-inflammatory effects of methylsulfonylmethane on lipopolysaccharide-induced inflammatory responses in murine macrophages. Center for Efficacy Assessment and Development of Functional Foods and Drugs, Hallym University, Korea. *Biol Pharm Bull.* 2009 Apr;32(4):651-6.
- Ko WC, Shih CM, Leu IJ, Chen TT, Chang JP. Mechanisms of relaxant action of luteolin in isolated guinea pig trachea. Graduate Institute of Pharmacology, College of Medicine, Taipei Medical University, Taipei, Taiwan, ROC. *Planta Med.* 2005 May;71(5):406-11.
- Kowalczyk MC, Kowalczyk P, Tolstykh O, Hanausek M, Walaszek Z, Slaga TJ. Synergistic effects of combined phytochemicals and skin cancer prevention in SENCAR mice. Department of Pharmacology, University of Texas Health Science Center at San Antonio, 78229, USA. *Comment in: Cancer Prev Res (Phila).* 2010 Feb;3(2):132-5. *Cancer Prev Res (Phila).* 2010 Feb;3(2):170-8. Epub 2010 Jan 26.
- Lansdown AB, Mirastschijski U, Stubbs N, Scanlon E, Agren MS. Zinc in wound healing: theoretical, experimental, and clinical aspects. Imperial College Faculty of Medicine, Division of Investigative Sciences, Charing Cross Hospital, London, United Kingdom. *Wound Repair Regen.* 2007 Jan-Feb;15(1):2-16.
- Laurent C, Voisin P, Pouget JP. DNA damage in cultured skin microvascular endothelial cells exposed to gamma rays and treated by the combination pentoxifylline and alpha-tocopherol. Service de Radiobiologie et d'Epidémiologie, DRPH, Institut de Radioprotection et de Sécurité Nucléaire, Fontenayaux-Roses, France. *Int J Radiat Biol.* 2006 May;82(5):309-21.
- Lawson BR, Belkowski SM, Whitesides JF, et al. Immunomodulation of Murine Collagen-Induced Arthritis by N, N-Dimethylglycine and a Preparation of Perna canaliculus. In: Anonymous. 2007.
- Lawson GW, Sato A, Fairbanks LA, Lawson PT. Vitamin E as a treatment for ulcerative dermatitis in C57BL/6 mice and strains with a C57BL/6 background. Division of Laboratory Animal Medicine, David Geffen School of Medicine, University of California-Los Angeles, Los Angeles, California 90095-1718, USA. *Contemp Top Lab Anim Sci.* 2005 May;44(3):18-21.
- Lima CC, Pereira AP, Silva JR, Oliveira LS, Resck MC, Grechi CO, Bernardes MT, Olímpio FM, Santos AM, Incerpi EK, Garcia JA. Ascorbic acid for the healing of skin wounds in rats. Universidade José do Rosário Vellano, Alfenas,



VetNova

CUTANIA® Chews

CUTANIA® MINI Chews



Next Generation Dermatological Supplement for Long Term Management of Hair and Skin Conditions in Dogs and Cats.

Data Sheet

MG, Brazil. Braz J Biol. 2009 Nov;69(4):1195-201.

- Liu J, Wan Y, Zhao Z, Chen H. Determination of the content of rosmarinic acid by HPLC and analytical comparison of volatile constituents by GC-MS in different parts of *Perilla frutescens* (L.). *Britt. Chem Cent J.* 2013 Apr 1;7(1):61
- Livine S, Myhre G, Smith G, et al. Effect of a nutritional supplement containing N,N-dimethylglycine (DMG) on the racing standard bred. In: Anonymous. 1982;4.
- Mahabir S, Frenkel K, Brady MS, Coit D, Leibes L, Karkoszka J, Berwick M. Randomized, placebo-controlled pilot trial of the effects of alpha-tocopherol supplementation on levels of autoantibodies against 5-hydroxymethyl-2-deoxyuridine in melanoma patients. Cancer Prevention Studies Branch, Center for Cancer Research, National Cancer Institute, Bethesda, Maryland 20892-8314, USA. *Melanoma Res.* 2004 Feb;14(1):49-56.
- Makino T, Furuta Y, Wakushima H, et al. Anti-allergic effect of *Perilla frutescens* and its active constituents. *Phytother Res.* 2003;17:240-3.
- Makino T, Furuta A, Fujii H, et al. Effect of oral treatment of *Perilla frutescens* and its constituents on type-I allergy in mice. *Biol Pharm Bull.* 2001;24:1206-9.
- Man J, Rao J, Goldman M. A double-blind, comparative study of nonanimal-stabilized hyaluronic acid versus human collagen for tissue augmentation of the dorsal hands. Department of Medicine, University of Alberta, Edmonton, Alberta, Canada. *Dermatol Surg.* 2008 Aug;34(8):1026-31. Epub 2008 Apr 22.
- Mani S, Lawson JW. Partial fractionation of Perna and the effect of Perna and Dimethylglycine on immune cell function and melanoma cells. In: Anonymous. Charleston, SC: South Carolina Statewide Research Conference, 1999.
- Mani S, Whitesides JF, Lawson JW. Role of Perna and Dimethylglycine (DMG) in modulating cytokine response and their impact on melanoma cells. In: Anonymous. Chicago, IL: 99th General Meeting of the American Society for Microbiology, 1999.
- McCusker MM, Grant-Kels JM. Healing fats of the skin: the structural and immunologic roles of the omega-6 and omega-3 fatty acids. Department of Dermatology University of Connecticut Health Center, 263 Farmington Avenue, MC 6230, Farmington, CT 06030, USA. 2010 Jul-Aug;28(4):440-51.
- Moffitt P, Potter G. Venous actic acid levels in exercising horses fed N,N-Dimethylglycine. In: Anonymous. Lansing, Michigan: Proceedings of the 9th Equine Nutrition and Physiology Symposium, 1985;248.
- Musalmah M, Nizrana MY, Fairuz AH, NoorAini AH, Azian AL, Gapor MT, Wan Ngah WZ. Comparative effects of palm vitamin E and alpha-tocopherol on healing and wound tissue antioxidant enzyme levels in diabetic rats. Department of Biochemistry, Faculty of Medicine, University Kebangsaan Malaysia, Kuala Lumpur, Malaysia. *Lipids.* 2005 Jun;40(6):575-80.
- Nassiri-Asl M, Hosseinzadeh H. Review of the pharmacological effects of *Vitis vinifera* (Grape) and its bioactive compounds. Department of Pharmacology, Faculty of Medicine, Qazvin University of Medical Sciences, Qazvin, IR Iran. *Phytother Res.* 2009 Sep;23(9):1197-204.
- Neuman MG, Oruña L, Coto G, Lago G, Nanau R, Vincent M. Hyaluronic acid signals for repair in ethanol-induced apoptosis in skin cells in vitro. Department of Clinical Pharmacology and Toxicology, University of Toronto, Toronto, ON, Canada M5S 1L7. *Clin Biochem.* 2010 Jul;43(10-11):822-6. Epub 2010 Apr 14.
- Oh HA, Park CS, Ahn HJ, Park YS, Kim HM. Effect of *Perilla frutescens* var. *acuta* Kudo and rosmarinic acid on allergic inflammatory reactions. *Exp Biol Med (Maywood).* 2011 Jan; 236(1):99-106.
- Osakabe N, Takano H, Sanbongi C, et al. Anti-inflammatory and anti-allergic effect of rosmarinic acid (RA); inhibition of seasonal allergic rhinoconjunctivitis (SAR) and its mechanism. *Biofactors.* 2005;21:127-31.
- Paniello RC, Hayden RE, Bello SL. Improved survival of acute skin flaps with amino acids as free radical scavengers. Department of Otolaryngology, Washington University School of Medicine, St Louis, MO 63110. *Arch Otolaryngol Head Neck Surg.* 1988 Dec;114(12):1400-3.
- Park J, Halliday GM, Surjana D, Damian DL. Nicotinamide prevents ultraviolet radiation-induced cellular energy loss. Discipline of Dermatology, Bosch Institute, Sydney Cancer Centre at Royal Prince Alfred Hospital, University of Sydney, Camperdown, New South Wales, Australia. *Photochem Photobiol.* 2010 Jul-Aug;86(4):942-8. Epub 2010 May 13.
- Pavicic T, Steckmeier S, Kerscher M, Korting HC. Evidence-based cosmetics: concepts and applications in photoaging of the skin and xerosis. *Klinik und Poliklinik für Dermatologie und Allergologie, Klinikum der Universität München, Germany. Wien Klin Wochenschr.* 2009;121(13-14):431-9.
- Raffoul W, Far MS, Cayeux MC, Berger MM. Nutritional status and food intake in nine patients with chronic low-limb ulcers and pressure ulcers: importance of oral supplements. *Chirurgie Plastique et Reconstructive, CHUV, Lausanne, Switzerland. Nutrition.* 2006 Jan;22(1):82-8.
- Reap EA, Lawson JW. Stimulation of the immune response by dimethylglycine, a nontoxic metabolite. *J Lab Clin Med.* 1990 Apr;115(4):481-6.



VetNova

CUTANIA® Chews

CUTANIA® MINI Chews

Next Generation Dermatological Supplement for Long Term Management of Hair and Skin Conditions in Dogs and Cats.



Data Sheet

- Reichrath J, Lehmann B, Carlberg C, Varani J, Zouboulis CC. Vitamins as hormones. Department of Dermatology, The Saarland University Hospital, Homburg, Germany. *Horm Metab Res.* 2007 Feb;39(2):71-84.
- Sanbongi C, Takano H, Osakabe N, Sasa N, Natsume M, Yanagisawa R, Inoue KI, Sadakane K, Ichinose T, Yoshikawa T. Rosmarinic acid in perilla extract inhibits allergic inflammation induced by mite allergen, in a mouse model. Health and Bioscience Laboratories, Meiji Seika Kaisha, Ltd, Sakado, Saitama, Japan. *Clin Exp Allergy.* 2004 Jun;34(6):971-7.
- Sharma SD, Katiyar SK. Dietary grape seed proanthocyanidins inhibit UVB-induced cyclooxygenase-2 expression and other inflammatory mediators in UVB-exposed skin and skin tumors of SKH-1 hairless mice. Department of Dermatology, University of Alabama, 1670 University Boulevard, Volker Hall 557, Birmingham, Alabama 35294, USA. *Pharm Res.* 2010 Jun;27(6):1092-102. Epub 2010 Feb 9.
- Schols JM, Heyman H, Meijer EP. Nutritional support in the treatment and prevention of pressure ulcers: an overview of studies with an arginine enriched oral nutritional supplement. Department of General Practice, Nursing Home Medicine, University Maastricht, P.O. Box 616, Maastricht 6200 MD. The Netherlands. *J Tissue Viability.* 2009 Aug;18(3):72-9. Epub 2009 May 8.
- Shin TY, Kim SH, Kim SH, Kim YK, Park HJ, Chae BS, Jung HJ, Kim HM. Inhibitory effect of mast cell-mediated immediate-type allergic reactions in rats by *Perilla frutescens*. College of Pharmacy, Woosuk University, Chonju, Chonbuk, South Korea. 2000 Aug;22(3):489-500.
- Schwartz JR, Marsh RG, Draelos ZD. Zinc and skin health: overview of physiology and pharmacology. Beauty Care Product Development, The Procter & Gamble Company, Cincinnati, Ohio 45251, USA. *Dermatol Surg.* 2005 Jul;31(7 Pt 2):837-47; discussion 847.
- Surjana D, Halliday GM, Damian DL. Role of nicotinamide in DNA damage, mutagenesis, and DNA repair. Discipline of Dermatology, Sydney Cancer Centre, Bosch Institute, University of Sydney at Royal Prince Alfred Hospital, Camperdown, Sydney, NSW 2006, Australia. *J Nucleic Acids.* 2010 Jul 25;2010. pii: 157591.
- Takano H, Osakabe N, Sanbongi C, et al. Extract of *Perilla frutescens* enriched for rosmarinic acid, a polyphenolic phytochemical, inhibits seasonal allergic rhinoconjunctivitis in humans. *Exp Biol Med (Maywood)* . 2004 Mar;229:247-54.
- Toxicological Research, Food and Drug Administration, Jefferson, Arkansas 72079, USA. *Vitam Horm.* 2007;75:223-56.
- Trookman NS, Rizer RL, Ford R, Mehta R, Gotz V J. Clinical assessment of a combination lip treatment to restore moisturization and fullness. *Clin Aesthet Dermatol* . 2009 Dec;2(12):44-8.
- Weiss RC. Immunologic responses in healthy random-source cats fed N,N-dimethylglycine-supplemented diets. *Am J Vet Res.* 1992 May;53(5):829-33.
- Wu G, Bazer FW, Burghardt RC, Johnson GA, Kim SW, Knabe DA, Li P, Li X, McKnight JR, Satterfield MC, Spencer TE. Proline and hydroxyproline metabolism: implications for animal and human nutrition. Department of Animal Science and Faculty of Nutrition, Texas A&M University, College Station, TX, 77843, USA. *Amino Acids.* 2010 Aug 10.
- Zouboulis CC. Retinoids which dermatological indications will benefit in the near future? Department of Dermatology, University Medical Center Benjamin Franklin, The Free University of Berlin, Berlin, Germany. *Skin Pharmacol Appl Skin Physiol.* 2001 Sep-Oct;14(5):303-15.

If you are interested in any of the articles listed, please do not hesitate to request them through the following contacts: vetnova@vetnova.net, +34 918 440 273, or your VetNova or Distributor Sales Representative.



VetNova

T: +34 918 440 273 · vetnova@vetnova.net · www.vetnova.net

VN-PUB-0089EN.0616